

Analysis on diurnal global geomagnetic variability under quiet-time conditions

Virginia Klausner^{1,*}

DGE/CEA/National Institute for Space Research - phone: +55 12 32086819/fax: +55 12 32086810 - INPE 12227-010 São José dos Campos, SP, Brazil.

Margarete Oliveira Domingues

LAC/CTE/National Institute for Space Research - phone: +55 12 32086542/fax: +55 12 32086375 - INPE 12227-010 São José dos Campos, SP, Brazil

Odim Mendes Jr

DGE/CEA/National Institute for Space Research - phone: +55 12 32087854/fax: +55 12 32086810 - INPE 12227-010 São José dos Campos, SP, Brazil

Andres Reinaldo Rodriguez Papa²

National Observatory - phone: +55 21 35049142/fax: +55 21 25807081 - ON 20921-400, RJ, Brazil

Peter Frick

Laboratory of Physical Hydrodynamics - phone: +7 342 2378322/fax: +7 342 2378487 - Institute of Continuous Media Mechanics, Perm, Russia

Abstract

This paper describes a methodology (or treatment) to establish a represen-

*Corresponding author

Email addresses: `virginia@dge.inpe.br` (Virginia Klausner), `mo.domingues@lac.inpe.br` (Margarete Oliveira Domingues), `odim@dge.inpe.br` (Odim Mendes Jr), `papa@on.br` (Andres Reinaldo Rodriguez Papa), `frick@icmm.ru` (Peter Frick)

¹National Observatory - ON 20921-400, RJ, Brazil.

²State University of Rio de Janeiro - UERJ 20550-900, RJ, Brazil

tative signal of the global magnetic diurnal variation based on a spatial distribution in both longitude and latitude of a set of magnetic stations as well as their magnetic behavior on a time basis. For that, we apply the Principal Component Analysis (PCA) technique implemented using gapped wavelet transform and wavelet correlation. The continuous gapped wavelet and the wavelet correlation techniques were used to describe the features of the magnetic variations at Vassouras (Brazil) and other 12 magnetic stations spread around the terrestrial globe. The aim of this paper is to reconstruct the original geomagnetic data series of the H-component taking into account only the diurnal variations with periods of 24 hours on geomagnetically quiet days. With the developed work, we advance a proposal to reconstruct the baseline for the quiet day variations (Sq) from the PCA using the correlation wavelet method to determine the global variation of PCA first mode. The results showed that this goal was reached and encourage other uses of this approach to different kinds of analysis.

Keywords: Magnetogram data, H-component, Quiet days, Principal Component Wavelet analysis.

1. Introduction

A substantial part of the energy carried by the solar wind can be transferred into the terrestrial magnetosphere and it is associated with the passage of southward directed interplanetary magnetic fields, B_s , by the Earth for sufficiently long intervals of time. Gonzalez et al. (1994) discussed the energy transfer process as a conversion of the directed mechanical energy from the solar wind into magnetic energy stored in the magnetotail of Earth's magne-

8 tosphere and its reconversion into thermal mechanical energy in the plasma
9 sheet, auroral particles, ring current, and Joule heating of the ionosphere.

10 The increase on the solar wind pressure is responsible for the energy in-
11 jections and induces global effects in the magnetosphere called geomagnetic
12 storms. The characteristic signature of geomagnetic storms can be described
13 as a depression on the horizontal component of the Earth’s magnetic field
14 measured at low and middle latitude ground stations. The decrease in the
15 magnetic horizontal field component is due to an enhancement of the trapped
16 magnetospheric particle population, consequently an enhanced ring of cur-
17 rent. This perturbation of the H-component could last from several hours to
18 several days (as described by Kamide et al., 1998).

19 The geomagnetic storms can consist of four phases: sudden commence-
20 ment, initial phase, main phase and recovery phase. The sudden commence-
21 ment when it exists, corresponds to the moment when the initial impact of
22 the increased solar wind pressure over the magnetopause occurs. The ini-
23 tial phase at ground appears as a rapid increase on the H-component over
24 less than 1 h almost simultaneously worldwide. The main phase of the ge-
25 omagnetic storm lasts a few hours and is characterized by a decrease in the
26 H-component. The recovery time corresponds to the gradual increase of the
27 H-component value to its average level. A detailed description of the mor-
28 phology of magnetic storms is, for instance, in Gonzalez et al. (1994).

29 The intensity of the geomagnetic disturbance in each day is described
30 by indices. The indices are very useful to provide the global diagnostic of
31 the degree of disturbance level. There are different indices that can be used
32 depending on the character and the latitude influences in focus. Considering

33 only the main latitudinal contributions, the ring current dominates at low
 34 and middle latitudes and an auroral ionospheric current systems dominates
 35 at higher latitudes (Mendes et al., 2005a). Kp, AE and Dst and their deriva-
 36 tions are the most used geomagnetic indices. The Kp index is obtained from
 37 the H-component and it is divided in ten levels from 0 to 9 corresponding to
 38 the mean value of the disturbance levels within 3-h intervals observed at 13
 39 subauroral magnetic stations (see Bartels, 1957). However, the K index is
 40 the most difficult to be physically interpreted due to its variations be caused
 41 by any geophysical current system including magnetopause currents, field-
 42 aligned currents, and the auroral electrojets (Gonzalez et al., 1994). The
 43 minutely AE index (sometimes 2.5 minute interval) is also obtained by the
 44 H-component measured from magnetic stations (5 to 11 in number) located
 45 at auroral zones and widely distributed in longitude. The AE index provides
 46 a measure of the overall horizontal auroral oval current strength. The index
 47 most used in low and mid-latitudes is the Dst index. It represents the varia-
 48 tions of the H-component due to changes of the ring current and is calculated
 49 every hour.

50 The Dst index is described as a measure of the worldwide derivation
 51 of the H-component at mid-latitude ground stations from their quiet days
 52 values. At mid-latitude, the H-component is a function of the magnetopause
 53 currents, the ring current and tail currents. Burton et al. (1975) calculated
 54 the Dst index as a average of the records from N mid-latitude magnetic
 55 stations following,

$$Dst = \frac{1}{N} \sum_{i=1}^N H_{disturbed} - H_{quiet} = \overline{\Delta H} \quad (1)$$

56 where $\overline{\Delta H}$ is a local time H average, $H_{disturbed}$ is the H-component mea-
 57 sured at disturbed days and H_{quiet} , on quiet days.

58 Other contributions beyond the ring current could be extracted or elim-
 59 inated with the idea presented by Burton et al. (1975). Those authors de-
 60 scribed the evolution of the ring current by a simple first order differential
 61 equation,

$$\frac{d Dst^*}{dt} = Q(t) - a Dst^*, \quad (2)$$

62 where $Dst^* = Dst - b Pdyn^{\frac{1}{2}} + c$. The contribution of the magnetopause
 63 currents to H is proportional to the square root of the solar wind dynamic
 64 pressure ($Pdyn$), Q represents the injection of particles to the ring current,
 65 $a Dst^*$ represents the loss of particles with an e-folding time $\frac{1}{a}$ and the con-
 66 stant terms a , b and c are determine by the quiet days values of the magne-
 67 topause and ring currents.

68 The Dst index is available on the Kyoto World Data Center at [http://](http://wdc.kugi.kyoto-u.ac.jp/dstdir/index.html)
 69 wdc.kugi.kyoto-u.ac.jp/dstdir/index.html. It is traditionally calculated from
 70 four magnetic observatories: Hermanus, Kakioka, Honolulu, and San Juan.
 71 These observatories are located at latitudes below 40° which are sufficiently
 72 distant from the auroral electrojets. The derivation of the Dst index corre-
 73 sponds to three main steps: the removal of the secular variation, the elimi-
 74 nation of the Sq variation and the calculation of the hourly equatorial Dst
 75 Index (see <http://wdc.kugi.kyoto-u.ac.jp/dstdir/dst2/onDstindex.html>).

76 The traditional method of calculating the baseline for the quiet day vari-
 77 ations uses the five quietest day for each month for each magnetic observa-
 78 tory. In this work, we propose a way to deal with Sq variations by suggesting

79 a method using Principal Component with the wavelet correlation matrix.
80 This method eliminates the disturbed days using a multiscale process. Also,
81 we developed an algorithm for extracting the solar quiet variations recorded
82 in the magnetic stations time series, in order words, a way of estimation of
83 the quiet-time baseline. To accomplish this task, we separate the solar di-
84 urnal variations using hourly data of the H-component using the technique
85 (described in Klausner et al., 2011a). Afterward we applied the principal
86 component wavelet analysis to identify the global patterns in the solar diur-
87 nal variations.

88 The rest of the paper is organized as follows: Section 2 is devoted to
89 explain the main issues of the Dst index calculation procedure. In Section 3
90 the analyzed period and data are presented. Section 4 describes the principal
91 component analysis and it is devoted to introduce the suggested method of
92 principal component analysis (PCA) using gapped wavelet transform and
93 wavelet correlation. It also establishes the identification of the disturbed
94 days. The results are discussed in Section 5, and finally, Section 6 brings the
95 conclusions of this work.

96 **2. The Dst index calculation procedure**

97 Recently, reconstructing the Dst and removing the quiet-time baseline
98 have been a motivation of several works (Karinen and Mursula, 2005; Mendes et al.,
99 2005a; Karinen and Mursula, 2006; Mursula et al., 2008; Love and Gannon,
100 2009; Klausner et al., 2011a). Despite of these disagreements in the Dst con-
101 struction, today, the Dst index remains an important tool in the space weather
102 analysis.

103 Karinen and Mursula (2005) reconstructed the Dst index (Dxt) follow-
104 ing the original formula presented at [http://wdc.kugi.kyoto-u.ac.jp /dst-](http://wdc.kugi.kyoto-u.ac.jp/dst-dir/dst2/onDstindex.html)
105 [dir/dst2/onDstindex.html](http://wdc.kugi.kyoto-u.ac.jp/dst-dir/dst2/onDstindex.html). However, they encountered a few issues as: the
106 availability and the data quality , some shifts in the baseline level of the
107 H-component, erroneous data points and some data gaps at all/some mag-
108 netic observatories. The Dst could not be fully reproduced using the original
109 formula because the inadequate information above effects the treatment of
110 the related issues and therefore remains partly unscientific as described by
111 Karinen and Mursula (2005).

112 A new corrected and extended version (Dcx) of the Dxt index was pro-
113 posed by Karinen and Mursula (2006). They corrected the Dst index for the
114 excessive seasonal varying quiet-time level which was unrelated to magnetic
115 storms as previously discussed in Karinen and Mursula (2005). They also
116 showed that the considerable amount of quiet-time variation is included in
117 the Dxt index but none in the Dcx index.

118 Another issue related to the derivation of the Dst index is that no treat-
119 ment is made to normalize the different latitudinal location of the magnetic
120 observatories. Mursula et al. (2008) suggested the normalization of the mag-
121 netic disturbances at the four Dst stations with different latitudes by the co-
122 sine of the geomagnetic latitude of the respective station. If no correction is
123 made, they showed for the lowest geomagnetic station, Honolulu, the largest
124 deviations and for the highest station, Hermanus, the lowest deviations of
125 the four station. The standard deviations reflect the annually averaged ef-
126 fect of the (mainly ring current related) disturbances at each station, (see
127 Mursula et al., 2008, for more details).

128 Mendes et al. (2005a) evaluated the effect of using more than four mag-
129 netic stations and shorter time intervals to calculate the Dst index. The
130 obtained Dst index profiles using 12, 6 or 4 magnetic station did not show
131 significant discrepancies and the best agreement with the standart Dst was
132 obtained using magnetic stations located at latitudes lower than 35° in both
133 hemispheres.

134 Although, the increase of symmetrically world-wide distributed magnetic
135 stations did not effect significantly the Dst index, the longitudinal asymme-
136 tries of the ring current contributes for the average disturbances of the Four
137 Dst stations be systematically different. Mursula et al. (2010) using an ex-
138 tended network of 17 stations, demonstrated that the local disturbances are
139 ordered according to the station’s geographic longitude, where the western-
140 most station (Honolulu) presented the largest disturbances and contributions
141 to Dst index and the easternmost (Kakioka) the smallest.

142 Klausner et al. (2011a) studied the characteristics of the Sq variations at
143 a Brazilian station and compared to the features from other magnetic sta-
144 tions to better understand the dynamics of the diurnal variations involved in
145 the monitoring of the Earth’s magnetic field. They used gapped wavelet anal-
146 ysis and the wavelet cross-correlation technique to verify the latitudinal and
147 longitudinal dependence of the diurnal variations. As previously mentioned
148 by Mursula et al. (2010), Klausner et al. (2011a) also verified that magnetic
149 stations located at lower latitudes and westernmost (Honolulu and San Juan)
150 presented larger correlation to Vassouras than the easternmost stations (as
151 Kakioka).

152 Some important aspects for the construction of the Dst index as described

by Love and Gannon (2009) are: the utilization of the original data, the inspection in time and frequency domains (removal of diurnal variation) and the consideration of the distinction between stationary and non-stationary time series ingredients which applies to the secular variation. Also as mentioned by Love and Gannon (2009), some patterns of the global magnetic disturbance field are well understood and some are not which means that there is still a lot to learn about the magnetosphere, magnetic storm and Earth-Sun relationship.

3. Magnetic Data

In this paper, we use ground magnetic measurements to estimate the quiet-time baseline. We select the four magnetic observatories used to calculate the Dst index: Hermanus (HER), Kakioka (KAK), Honolulu (HON), and San Juan (SJG), plus other 9 different magnetic observatories reasonably homogeneously distributed world wide. One of these nine chosen stations is Vassouras (VSS) located under the South Atlantic Magnetic Anomaly (minimum of the geomagnetic field intensity). The geomagnetic data use in this work relied on data collections provided by the INTERMAGNET programme (<http://www.intermagnet.org>).

The distribution of the magnetic stations, with their IAGA codes, is given in Fig. 1. The corresponding codes and locations are given in Table 1. These selection of magnetic stations correspond to the same selection used in a previous work of (Klausner et al., 2011a) for the same reasons (exclusion of the major influence of the auroral and equatorial electrojets). In this work, we only use the data interval corresponding to the year 2007. We also ap-

177 ply the same methodology to identify geomagnetically quiet days used by
 178 Klausner et al. (2011a). We consider quiet days, only those days in which
 179 the Kp index is not higher than 3+.

180 As at low latitudes the horizontal component (H) is mostly affected by
 181 the intensity of the ring current, we decided to use only the hourly mean
 182 value series of this component. The magnetic stations present available data
 183 in Cartesian components (XYZ system). The Conversion to horizontal-polar
 184 components (HDZ system) is very simple (see Campbell, 1997, for more
 185 details). The system's conversion was performed in all the chosen magnetic
 186 stations.




Figure 1: Geographical localization of the stations used in this work and their respective IAGA code.

187 4. Methodology

188 The method used in this study is based on the principal component analy-
 189 sis (PCA) using gapped wavelet transform and wavelet correlation to charac-

Table 1: INTERMAGNET network of geomagnetic stations used in this study.

Station	Geographic coord.		Geomagnetic coord.	
	Lat.($^{\circ}$)	Long.($^{\circ}$)	Lat.($^{\circ}$)	Long.($^{\circ}$)
AMS	-37.83	77.56	-46.07	144.94
ASP	-23.76	133.88	-32.50	-151.45
BEL	51.83	20.80	50.05	105.18
BMT	40.30	116.20	30.22	-172.55
BOU	40.13	-105.23	48.05	-38.67
CLF	48.02	2.26	49.56	85.72
CMO	64.87	-147.86	65.36	-97.23
EYR	-43.42	172.35	-46.79	-106.06
HER	-34.41	19.23	-33.89	84.68
HON	21.32	-158.00	21.59	-89.70
KAK	36.23	140.18	27.46	-150.78
SJG	18.12	-66.15	27.93	6.53
VSS	-22.40	-43.65	-13.43	27.06

Source: <http://wdc.kugi.kyoto-u.ac.jp/igrf/gggm/index.html> (2010)

190 terize the global diurnal variation behavior. To identify periods of magnetic
 191 disturbance, we use the discrete wavelet transform. This technique is em-
 192 ployed to analyze the removal of disturbed days from the magnetograms, and
 193 consequently, from the reconstructed Sq signal. Also in this section, a com-
 194 bined methodology using the PCA and gapped wavelet transform is briefly
 195 described. Following, we present an identification method to distinguish the
 196 disturbed days using discrete wavelet coefficients.

198 Among the several available methods of analysis, PCA is a particularly
199 useful tool in studying large quantities of multi-variate data. PCA is used
200 to decompose a time-series into its orthogonal component modes, the first of
201 which can be used to describe the dominant patterns of variance in the time
202 series Murray (1984). The PCA is able also to reduce the original data set
203 of two or more observed variables by identifying the significant information
204 from the data. Principal Components (PCs) are derived as the eigenvectors of
205 the correlation matrix between the variables. Their forms depend directly on
206 the interrelationships existing within the data itself. The first PC is a linear
207 combination of the original variables, which when used as a linear predictor
208 of these variables, explains the largest fraction of the total variance. The
209 second, third PC, etc., explain the largest parts of the remaining variance
210 Murray (1984).

211 As explained by Yamada (2002), the interpretation of the eigenvectors
212 and the eigenvalues can be described as follow, the eigenvectors are the nor-
213 malized orthogonal basis in phase space, and also, the set of vectors of the
214 new coordinate system in the space, different from the coordinate system of
215 the original variables; the eigenvalues are the corresponding variance of the
216 distribution of the projections in the new basis.

217 In order to isolate the global contributions of each PCs mode, we applied
218 PCA using the wavelet correlation matrix computed by gapped wavelet trans-
219 form. This wavelet correlation matrix was introduced in Nesme-Ribes et al.
220 (1995). We joined the properties of the PCA, which are the compression of
221 large databases and the simplification by the PCs modes, and properties of

222 the wavelet correlation matrix, which is the correlation at a given scale, a ,
223 in this case, the scale corresponded to the pseudo-period of 24 hours.

224 *Identification of magnetic disturbance*

225 The wavelet analysis has the following propriety: the larger amplitudes of
226 the wavelet coefficients are associated with locally abrupt signal changes or
227 “details” of higher frequency. In the work of Mendes et al. (2005b) and the
228 following work of Mendes da Costa et al. (2011), a method for the detection
229 of the transition region and the exactly location of this discontinuities due
230 to geomagnetic storms was implemented. In these cases, the highest ampli-
231 tudes of the wavelet coefficients indicate the singularities on the geomagnetic
232 signal in association with the disturbed periods. On the other hand, when
233 the magnetosphere is under quiet conditions for the geomagnetic signal, the
234 wavelet coefficients show very small amplitudes. In this work, we applied this
235 methodology with Daubechies orthogonal wavelet function of order 2 on the
236 one minute time resolution with the pseudo-periods of the first three levels
237 of 3, 6 and 12 minutes.

238 **5. Results and Discussion**

239 In this section, we will present the results of reconstructed baseline for the
240 global quiet days variation using PCA technique implemented with gapped
241 wavelet transform and wavelet correlation. Also, we will apply DWT to eval-
242 uate the day-by-day level of geomagnetic disturbance using KAK magnetic
243 station as reference.

244 Fig. 2 shows an example of the geomagnetic behavior presented at the
245 June, 2007 magnetogram of Kakioka using continuous gapped wavelet trans-




Figure 2: The GWT analysis of KAK magnetogram. At top, it shows the H-component of KAK at June, 2007 used for the wavelet analysis, at bottom left, the scalogram using Morlet wavelet, logarithmic scaled representing $\log 2(|W(a,b)|)$, and at bottom right, the global wavelet spectrum.

form (GWT). The GWT can be used in the analysis of non-stationary signal to obtain information on the frequency or scale variations and to detect its structures localization in time and/or in space (see Klausner et al., 2011a, for more details). It is possible to analyze a signal in a time-scale plane, called so the wavelet scalogram. In analogy with the Fourier analysis, the square modulus of the wavelet coefficient, $|W(a,b)|^2$, is used to provide the energy distribution in the time-scale plane. In the GWT analysis, we can also explore the central frequencies of the time series through the global wavelet spectrum which is the variance average at each scale over the whole time series, to compare the spectral power at different scales. This figure shows the H-component (top), the wavelet square modulus (bottom left) and the global wavelet spectrum (total energy in each scale – bottom right). In the

scalogram, areas of stronger wavelet power are shown in dark red on a plot of time (horizontally) and time scale (vertically). The areas of low wavelet power are shown in dark blue.

In Fig. 2, it is possible to notice peaks of wavelet power on the scalogram at the time scale corresponding to 8 to 16 minutes of period. These periods are associated to PC5 pulsations during disturbed periods. Also, it is possible to notice a maximum of wavelet power at the time scale corresponding to harmonic periods of the 24 hours such as 6, 8, 12 hours. Those periods are related to the diurnal variations.

The GWT technique is able to analyze all the informations present on the magnetograms. It is an auxiliary tool to localize on time/space the PC1–PC5 pulsations (Saito, 1969). However, the scalogram provides a very redundancy information which difficult the analysis of each decoupling phenomenon. For that reason, we preferred to use DWT to evaluate the day-by-day level of geomagnetic disturbance.

Fig. 3 is composed of two graphs, the first one presents the reconstructed line of the Sq variation where the first 10 geomagnetically quietest days of each month are highlighted in blue and the 5 most disturbed days in red and the second one presents the discrete wavelet analysis of the geomagnetic horizontal component obtained at Kakioka station, Japan.

Using our criteria of removing disturbed days, we consider as gaps of 3rd, 14th, 21th and 29th day. In our case, the gapped wavelet technique is very helpful because it reduces two effects: the presence of gaps and the boundary effects due to the finite length of the data, for more details see Klausner et al. (2011a).




Figure 3: The amplitude variation of the reconstructed Sq signal with the highlighted 10 geomagnetically quietest days (blue) and 5 most disturbed days (red), the H-component average variation for KAK obtained at June, 2007 and the square root wavelet coefficients amplitudes $d1$, $d2$ and $d3$ with the pseudo-periods of 3, 6 and 12 minutes.

283 The first graph shows the amplitude range between -5 and 5 nT and
 284 presents a complex pattern. It is possible to notice that the larger amplitudes
 285 of the reconstructed Sq signal correspond to the periods between the days
 286 8–10, 13–15, 21–23 and 29–30. These periods correspond to the disturbed
 287 days.

288 Table 5 shows these 10 quietest days and 5 most disturbed days of each
 289 month set by the *GeoForschungsZentrum (GFZ) Potsdam* through the anal-
 290 ysis of the Kp index that are highlighted on the second graph. The year
 291 of 2007 is a representative year of minimum solar activity and it is used in
 292 this analysis due to have less disturbed periods (see our considerations in
 293 Section 3). By analyzing these highlighted days, we expect to find out if
 294 there is a correlation between the days classified as quiet days and a small

295 Sq amplitude variation.

Table 2: The first 10 geomagnetically quietest days and first 5 most disturbed days set by the *GeoForschungsZentrum (GFZ) Potsdam*

Month/year	10 quietest days										5 most disturbed days				
	q1	q2	q3	q4	q5	q6	q7	q8	q9	q10	d1	d2	d3	d4	d5
Mar	20	21	3	19	9	29	22	18	31	4	13	24	6	7	14
Jun	5	12	6	7	11	26	25	20	30	28	14	21	22	3	29
Sep	13	9	10	11	12	16	17	19	26	15	29	2	28	23	27
Dec	3	8	4	25	7	26	2	29	15	6	18	17	11	20	21

SOURCE: <http://wdc.kugi.kyoto-u.ac.jp/qddays/index.html>.

296 The second graph shows the discrete wavelet analysis applied to geo-
 297 magnetic minutely signal from KAK using Daubechies orthogonal wavelet
 298 family 2. From top to bottom in this graph, the H-component of the ge-
 299 omagnetic field and the first three levels of the square wavelet coefficients
 300 denoted by d1, d2 and d3. This analysis uses the methodology developed
 301 by Mendes et al. (2005b), and posteriorly applied by Mendes da Costa et al.
 302 (2011) and Klausner et al. (2011b).

303 In order to facilitate the evaluation of the quiet periods obtained by the
 304 discrete wavelet analysis applied to geomagnetic signal from KAK, we also
 305 developed a methodology (effectiveness wavelet coefficients (EWC)) to in-
 306 terpret the results shown in Fig. 4. The EWC corresponds to the weighted
 307 geometric mean of the square wavelet coefficients per hour. It is accom-
 308 plished by weighting the square wavelet coefficients means in each level of
 309 decomposition as following

$$EWC = \frac{4 \sum_{i=1}^N d1 + 2 \sum_{i=1}^N d2 + \sum_{i=1}^N d3}{7}, \quad (3)$$

310 where N is equal to 60 because our time series has one minute resolution.




Figure 4: The comparative of the global Sq behavior and the effectiveness wavelet coefficients for the month of June, 2007.

Through Fig. 4, it is possible to compare the global Sq behavior (top graph) with the analysis of quiet and disturbed days obtained by one representative magnetic station of medium/low latitudes (KAK – bottom graph) in order to verify situations in which the global Sq behavior presents less or more variability. This analysis allows us to validate the quietest days, and evaluate the most disturbed days in order to establish a reliable method of global Sq analysis obtained from medium/low latitude magnetic stations influenced only by the ionosphere.

In Fig. 4, the global Sq behavior (top graph) shows larger amplitudes during the periods between the days 8–10, 12–16, 22–23 and 29–30. Most of these periods correspond to the days where the EWCs have an increase of their values as shown KAK analysis (bottom graph). The increase of the EWCs values occurs during the periods between the days 1–4, 8–10, 13–18,

21–24 and 29–30. The EWCs can help us also to interpret the results obtained in each day, and can help us to evaluate the quietest and most disturbed days measure by the selected magnetic station of medium/low latitudes, KAK, as shown in Table 5.

Table 3: The 10 geomagnetically quietest days and 5 most disturbed days obtained by the discrete wavelet analysis

Month/year	10 quietest days										5 most disturbed days				
	q1	q2	q3	q4	q5	q6	q7	q8	q9	q10	d1	d2	d3	d4	d5
Mar	3	10	20	9	21	8	2	19	22	4	13	25	12	15	24
Jun	11	12	6	25	5	26	7	28	19	27	21	2	14	22	29
Sep	12	11	10	9	17	18	16	13	26	19	27	22	29	2	20
Dec	25	7	3	8	1	24	4	6	16	28	17	18	11	20	10

The same methodology and analysis comparing the global Sq behavior and the EWCs from KAK is done for the month of March, September and December, 2007.

Fig. 5 shows the comparative of the global Sq behavior and the EWCs from KAK for the month of March, 2007. The amplitude range of global Sq signal is between -15 and 15 nT, and, it also shows a complex pattern. It is possible to notice that the larger amplitudes of the reconstructed Sq signal correspond to the periods between the days 6–7, 11–17 and 23–28. Once more, these periods correspond to the disturbed days. The increase of the EWCs values occurs during the periods between the days 1, 6–7, 11–17, 23–28 and 30–31. We can notice that the increase of the amplitude of the reconstructed Sq signal correspond to the increase of the EWCs magnitude.

The amplitude range for September, 2007, is between -15 and 15 nT and the larger amplitudes of the reconstructed Sq signal correspond to the periods between the days 1–7, 19–23 and 26–30, see Fig. 6. The increase of




Figure 5: The comparative of the global Sq behavior and the effectiveness wavelet coefficients for the month of March, 2007

the EWCs values occurs during the periods between the days 1–3, 20–24 and 27–30. Comparing these two analysis, the global and the EWCs, we verify that the KAK magnetic behavior represents well the increase of global Sq oscillations.

In Fig. 7, the amplitude range is between -10 and 10 nT and the larger amplitudes correspond to the periods between the days 15–23. Also, the increase of the EWCs values occur during the periods between the days 10–11 and 17–21. Once more, when we compare these two analysis, the global and the EWCs, we verify that the KAK magnetic behavior represents well the increase of global Sq oscillations.

We observe in Figs. 4, 5, 6 and 7, in most of the cases, the major amplitude fluctuations of the reconstructed Sq signal correspond to the most disturbed days and minor fluctuations, to the quietest days. However, the




Figure 6: The comparative of the global Sq behavior and the effectiveness wavelet coefficients for the month of September, 2007

356 diurnal global variability shows a complexity on the amplitude variation pat-
 357 tern even during geomagnetically quiet periods. Through this study we com-
 358 pare the amplitude variation of the reconstructed Sq signal to effectiveness
 359 wavelet coefficients obtained at KAK with the purpose of understanding the
 360 complexity of the diurnal global variability.

361 6. Conclusions

362 In this work, we suggest an alternative approach for the calculation of the
 363 Sq baseline using wavelet and PCA techniques. This new approach address
 364 some issues, such as, the availability and the quality of data, abrupt changes
 365 in the level of the H-component, erroneous points in the database and the
 366 presence of gaps in almost all the magnetic observatories. To fulfill this
 367 purpose, we reconstruct the Sq baseline using the wavelet correlation matrix




Figure 7: The comparative of the global Sq behavior and the effectiveness wavelet coefficients for the month of December, 2007

with scale of 24 hours (pseudo-period). The PCA/wavelet method uses the global variation of first PCA mode that also corresponds to phenomena with periods of 24 hours. This study shows that the largest amplitude oscillation of the reconstructed signal (Sq baseline) corresponded to the most disturbed days and the smaller oscillations to the quietest days. This result is consistent with the expected Sq variations.

7. Acknowledgments

V. Klausner wishes to thanks CAPES for the financial support of her PhD (CAPES – grants 465/2008) and her Postdoctoral research (FAPESP – 2011/20588-7). This work was supported by CNPq (grants 309017/2007-6, 486165/2006-0, 308680/2007-3, 478707/2003, 477819/2003-6, 382465/01-6), FAPESP (grants 2007/07723-7) and CAPES (grants 86/2010-29, 0880/08-6,

380 86/2010-29, 551006/2011-0, 17002/2012-8). Also, the authors would like to
381 thank the INTERMAGNET programme for the datasets used in this work.

382 **References**

383 Bartels, J., 1957. The technique of scaling indices k and q of geomagnetic
384 activity. *Annals of the International Geophysical Year* (4), 215–226.

385 Burton, R. K., McPherron, R. L., Russell, C. T., 1975. An empirical relation-
386 ship between interplanetary conditions and Dst. *Journal of Geophysical*
387 *Research*, 80 (1), pp. 4204–4214.

388 Campbell, W. H., 1997. *Introduction to Geomagnetic Fields*. Cambridge Uni-
389 versity Press, New York.

390 Domingues, M. O., Mendes, O. J., Mendes da Costa, A., 2005. Wavelet
391 techniques in atmospheric sciences. *Advances in Space Research* 35 (5),
392 831–842.

393 Gonzalez, W. D., Joselyn, J. A., Kamide, Y., Kroehl, H. W., Rostoker, G.,
394 Tsurutani, B. T., Vasyliunas, V. M., 1994. What is a geomagnetic storm?
395 *Journal of Geophysical Research*, 99 (A4), pp. 5771–5792.

396 Kamide, Y., Baumjohann, W., Daglis, I. A., Gonzalez, W. D., Grande, M.,
397 Joselyn, J. A., McPherron, R. L., Phillips, J. L., Reeves, E. G. D., Ros-
398 toker, G., Sharma, A. S., Singer, H. J., Tsurutani, B. T., and Vasyliunas,
399 V. M., 1998. Current understanding of magnetic storms: Storm-substorm
400 relationships. *Journal of Geophysical Research*, 103 (A8), pp. 17705–17713.

401 Karinen, A., Mursula, K., 2005. A new reconstruction of the Dst index for
402 1932-2002. *Annales Geophysicae*, 23, pp 475–485.

403 Karinen, A., Mursula, K., 2006. Correcting the Dst index: Consequences
404 for absolute level and correlations. *Journal of Geophysical Research*,
405 111 (A08207).

406 Klausner, V., Papa, A. R. R., Mendes, O., Domingues, M. O., Frick, P.,
407 2011a. Characteristics of solar diurnal variations: a case study based
408 on records from the ground magnetic observatory at Vassouras, Brazil.
409 arXiv:1108.4594 [physics.space-ph].

410 Klausner, V., Domingues, M. O., Mendes, O., Papa, A. R. R.,
411 2011b. Tsunami effects on the Z component of the geomagnetic field.
412 arXiv:1108.4893v1 [physics.space-ph]

413 Love, J. J., Gannon, J. L., 2009. Revised Dst and the epicycles of magnetic
414 disturbance: 19582007. *Annales Geophysicae* 27, pp. 3101–3131.

415 Mendes, O. J., Domingues, M. O., Mendes da Costa, A., Clúa de Gonza-
416 lez, A. L., 2005a. Wavelet analysis applied to magnetograms: Singular-
417 ity detections related to geomagnetic storms. *Journal of Atmospheric and*
418 *Solar-Terrestrial Physics*, 67, pp. 1827–1836.

419 Mendes, O. J., Mendes da Costa, A., Domingues, M. O., 2005b. Introduction
420 to planetary electrodynamics: a view of electric fields, currents and related
421 magnetic fields. *Advances in Space Research*, 35 (5), pp. 812-818.

422 Mendes da Costa, A., Domingues, M. O., Mendes, O., Brum, C. G. M.,

423 2011. Interplanetary medium condition effects in the south atlantic mag-
 424 netic anomaly: A case study. *Journal of Atmospheric and Solar-Terrestrial*
 425 *Physics*, 73 (11–12), pp. 1478–1491.

426 Murray, G. W.; Mueller, J. C.; Zwally, H. J., 1984 Matrix partitioning and
 427 EOF/Principal Components Analysis of Antarctic Sea Ice Brightness Tem-
 428 peratures. Greenbelt, Md. : National Aeronautics and Space Administra-
 429 tion, Goddard Space Flight Center, 1 v., NASA technical memorandum,
 430 83916.

431 Mursula, K., Holappa, L., Karinen, A., 2008. Correct normalization of the
 432 Dst index. *Astrophysics and Space Sciences Transactions*, 4 (2), pp.41–45.

433 Mursula, K., Holappa, L., Karinen, A., 2010. Uneven weighting of stations
 434 in the Dst index. *Journal of Atmospheric and Solar-Terrestrial Physics*,
 435 73 (2-3), pp. 316–322.

436 Nesme-Ribes, E., Frick, P., Sokoloff, D., Zakharov, V., Ribes, J. C.,
 437 Vigouroux, A., Laclare, F., 1995. Wavelet analysis of the maunder mini-
 438 mum as recorded in solar diameter data. *Comptes rendus de l’Académie*
 439 *des sciences. Série II, Mécanique, physique, chimie, astronomie* 321 (12),
 440 525–532.

441 Saito, T., 1969. Geomagnetic pulsations. *Space Science Reviews*, 10 (3), pp.
 442 319–412.

443 Yamada, K., 2002. 2-day, 3-day, and 5-6-day oscillations of the geomagnetic
 444 field detected by principal component analysis. *Earth Planets Space* 54
 445 (4), 379–392.